

850 W South Boulder Rd Ste 201 Louisville, CO 80027 303-665-6120 Westminster, CO 80234 303-214-4100

Before Endodontic Treatment

This procedure will be performed using local anesthesia. There are usually no restrictions after the procedure concerning driving or returning to work. A doctor is available for consultation at all times should a problem arise after your treatment.

Continue all medications for blood pressure, diabetes, thyroid problems and any other conditions as recommended by your physician. If there is a question, please call our office prior to your appointment.

If you have been advised by your physician or dentist to use antibiotic premedication because of mitral valve prolapsed (MVP), heart murmur, hip, knee, cardiac or other prosthesis, or if you have rheumatic heart disease, please make sure you are on the appropriate antibiotic on the day of your appointment. If there is a question, please call our office prior to your appointment.

If you can take ibuprofen (Advil) or naproxen sodium (Aleve), it does help reduce inflammation when taken pre-operatively. We recommend 2 tablets or either medication 2-4 hours before endodontic therapy.

General Instructions

Your tooth and surrounding gum tissue may be slightly tender for several days to several weeks as a result of manipulation during treatment and previous condition of your tooth. This tenderness is normal and is no cause for alarm. Do not chew food on the affected side until your endodontic therapy is completed and your tooth is covered with a protective restoration provided by your restorative dentist. You may continue you regular dental hygiene regimen. Discomfort may be alleviated by taking ibuprofen (Advil), aspirin, or acetaminophen (Tylenol) as directed. Should you experience discomfort that cannot be controlled with the above listed medications, or should swelling develop, please contact this office immediately. If you need to call after hours, please have your pharmacy number available.